

2009-2010 H1N1 Influenza Vaccine Dosage Chart (October 1, 2009)

Vaccine	Age Groups	Number of doses*	Dosage
Fluzone® (sanofi pasteur) Preservative Free 0.25 mL prefilled syringes	6 – 35 months	2	0.25 mL
Fluzone® (sanofi pasteur) Multi-dose vials	6 – 35 months	2	0.25 mL
	36 months and older	2 doses for people ages 9 and younger 1 dose for people ages 10 and older	0.5 mL
Fluzone® (sanofi pasteur) Preservative Free 0.5 mL vial and prefilled syringes	36 months and older	2 doses for people ages 9 and younger 1 dose for people ages 10 and older	0.5 mL
Fluvirin® (Novartis) Multi-dose vials	4 years and older	2 doses for people ages 9 and younger 1 dose for people ages 10 and older	0.5 mL
Fluvirin® (Novartis) Trace Preservative 0.5 mL prefilled syringes	4 years and older	2 doses for people ages 9 and younger 1 dose for people ages 10 and older	0.5 mL
Afluria® CSL Limited Preservative Free 0.5 mL prefilled syringes	18 years and older	1	0.5 mL
Afluria® CSL Limited Multi-dose vials	18 years and older	1	0.5 mL
FluLaval® GlaxoSmithKline	Not yet approved by the Food and Drug Administration		
FluMist®** (MedImmune) Preservative Free 0.2 mL Sprayers	2 through 49 years	2 doses for people ages 9 and younger 1 dose for people ages 10 and older	0.2 mL (0.1 mL per nostril)

*Most like three or four weeks apart (unknown as of October 1, 2009)

**H1N1 influenza vaccine may be administered at the same time as other vaccines, including seasonal influenza vaccine with one exception, if administering live intranasal influenza vaccine (Flumist®) for both H1N1 and seasonal doses, then the doses may not be administered on the same day, and must be separated by at least four weeks.

For more information about the above H1N1 influenza vaccines, including precautions and contraindications, please read the package insert.